ROSLINDALE PEDIATRIC ASSOCIATES, P.C.

PEDIATRICS AND ADOLESCENT MEDICINE

TYPICAL LAB & VACCINE SCHEDULE

WELL VISIT (Age)	LAB	VACCINE
BIRTH (Usually in Hospital)	none	HEPATITIS B #1 RSV Antibody (if applicable)
2 MONTHS	none	VAXELIS #1 (DTaP, IPV, Hib, Hep B) PREVNAR #1 (Pneumococcal) ROTATEQ #1 (Rotavirus)
4 MONTHS	none	VAXELIS #2, PREVNAR #2 ROTATEQ #2
6 MONTHS	none	VAXELIS #3, PREVNAR #3 ROTATEQ #3
9 MONTHS	FINGER STICK for Lead and Hemoglobin	none
12 MONTHS	none	MMR #1 (Measles, Mumps, Rubella) PREVNAR #4 VARIVAX #1 (Varicella)
15 MONTHS	none	VAQTA #1 (Hep A) PENTACEL (DTap #4, IPV #4, Hib #4)
18 MONTHS	FINGER STICK for Lead and Hemoglobin	none
2 YEARS	FINGER STICK for Lead and Hemoglobin	VAQTA #2
3 YEARS	FINGER STICK for Lead and Hemoglobin	none
4 YEARS	FINGER STICK for Lead and Hemoglobin (if high risk)	PROQUAD (MMR #2, Varicella #2) QUADRACEL (DTap #5, IPV #5)
5 to 8 YEARS	none	none
9 YEARS	none	GARDASIL #1 (HPV) GARDASIL #2 6 months later
10 YEARS	FINGER STICK for lipid panel	none
11 YEARS	none	ADACEL (Tdap) MENQUADFI #1 (Meningococcal)
12 to 15 YEARS	none	none
16 YEARS	Chlamydia screen (urine or swab)	MENQUADFI #2 BEXSERO (Men B) #1 BEXSERO #2 6 months later
17 YEARS	FINGER STICK for lipid panel Chlamydia screen (urine or swab)	none
18 to 20 YEARS	Chlamydia screen (urine or swab)	none
21 to 22 YEARS	Chlamydia screen (urine or swab)	ADACEL (Tdap)

Updated 4/11/2025

ROSLINDALE PEDIATRIC ASSOCIATES, P.C.
PEDIATRICS AND ADOLESCENT MEDICINE

- **NO STATE VACCINES for patients 19 years and older.**
- **ALL PATIENTS OLDER THAN 18 years need to use non-state vaccines.**
- **DURING THE FLU/COVID SEASON, at all visits 6 months old and up, FLU and/or COVID vaccines may be added if not yet vaccinated.**

Updated 4/11/2025